

Improved Constructions for Non-adaptive Threshold Group Testing

Mahdi Cheraghchi*

Abstract

The basic goal in combinatorial group testing is to identify a set of up to d defective items within a large population of size $n \gg d$ using a pooling strategy. Namely, the items can be grouped together in pools, and a single measurement would reveal whether there are one or more defectives in the pool. The threshold model is a generalization of this idea where a measurement returns positive if the number of defectives in the pool exceeds a fixed threshold u , negative if this number is below a fixed lower threshold $\ell \leq u$, and may behave arbitrarily otherwise. We study non-adaptive threshold group testing (in a possibly noisy setting) and show that, for this problem, $O(d^{g+2}(\log d) \log(n/d))$ measurements (where $g := u - \ell$ and u is any fixed constant) suffice to identify the defectives, and also present almost matching lower bounds. This significantly improves the previously known (non-constructive) upper bound $O(d^{u+1} \log(n/d))$. Moreover, we obtain a framework for explicit construction of measurement schemes using lossless condensers. The number of measurements resulting from this scheme is ideally bounded by $O(d^{g+3}(\log d) \log n)$. Using state-of-the-art constructions of lossless condensers, however, we come up with explicit testing schemes with $O(d^{g+3}(\log d) \text{quasipoly}(\log n))$ and $O(d^{g+3+\beta} \text{poly}(\log n))$ measurements, for arbitrary constant $\beta > 0$.

1 Introduction

Combinatorial group testing is a classical problem that deals with identification of sparse Boolean vectors using disjunctive queries. Suppose that among a large set of n items it is suspected that, for some *sparsity parameter* $d \ll n$, up to d items might be “defective”. In technical terms, defective items are known as *positives* and the rest are called *negatives*. In a *pooling strategy*, the items may be arbitrarily grouped in pools, and a single “measurement” reveals whether there is one or more positives within the chosen pool. The basic goal in group testing is to design the pools in such a way that the set of positives can be identified from a number of measurements that is substantially less than n .

Since its introduction in 1940’s [15], group testing and its variations have been extensively studied and have found surprisingly many applications in seemingly unrelated areas. In particular, we mention applications in molecular biology and DNA library screening (cf. [3, 19, 27, 29, 33, 38, 39] and the references therein), multi-access communication [37], data compression [23], pattern matching [11], streaming algorithms [12], software testing [2], compressed sensing [13], and secure key distribution [5], among others. We refer the reader to [16, 17] for an extensive review of the major results in this area.

*Department of Computer Science, University of Texas at Austin, USA. Email: mahdi@cs.utexas.edu. Part of work was done while the author was with the School of Computer and Communication Sciences, Ecole Polytechnique Fédérale de Lausanne (EPFL), Switzerland. Research was supported in part by the ERC Advanced investigator grant 228021 of A. Shokrollahi. A preliminary summary of this work appears (under the same title) in proceedings of the 37th International Colloquium on Automata, Languages and Programming (ICALP 2010), Bordeaux, France, July 2010 [10].

Formally, in classical group testing one aims to learn an unknown d -sparse¹ Boolean vector $(x_1, \dots, x_n) \in \{0, 1\}^n$ using a set of m measurements, where each measurement is defined by a subset of the coordinates $\mathcal{I} \subseteq [n]$ and outputs the logical “or” $\bigvee_{i \in \mathcal{I}} x_i$. The goal is then to design the measurements in such a way that all d sparse vectors become uniquely identifiable using as few number of measurements as possible.

A natural generalization of classical group testing (that we call *threshold testing*), introduced by Damaschke [14], considers the case where the measurement outcomes are determined by a *threshold predicate* instead of the logical or. Namely, this model is characterized by two integer parameters ℓ, u such that $0 < \ell \leq u$ (that are considered fixed constants), and each measurement outputs positive if the number of positives within the corresponding pool is at least u . On the other hand, if the number of positives is strictly less than² ℓ , the test returns negative, and otherwise the outcome can be arbitrary. In this view, classical group testing corresponds to the special case where $\ell = u = 1$. In addition to being of theoretical interest, the threshold model is interesting for applications, in particular in biology, where the measurements have reduced or unpredictable sensitivity or may depend on various factors that must be simultaneously present in the sample to result in a positive outcome.

The difference $g := u - \ell$ between the thresholds is known as the *gap* parameter. As shown by Damaschke [14], in threshold group testing identification of the set of positives is only possible when the number of positives is at least u . Moreover, regardless of the number of measurements, in general the set of positives can be only approximately identified within up to g false positives and g false negatives (thus, unique identification can only be guaranteed when $\ell = u$). Additionally, Damaschke constructed a scheme for identification of the positives in the threshold model. For the gap-free case where $g = 0$, the number of measurements in this scheme is $O(d \log n)$, which is nearly optimal (within constant factors). However, when $g > 0$, the number of measurements becomes $O(dn^b + d^u)$, for an arbitrary constant $b > 0$, if up to $g + (u - 1)/b$ misclassifications are allowed.

A drawback of the scheme presented by Damaschke is that the measurements are *adaptive*; i.e., the group chosen by each measurement can depend on the outcomes of the previous ones. For numerous applications (in particular, in molecular biology), adaptive measurements are infeasible and must be avoided. In a *non-adaptive* setting, all measurements must be specified before their outcomes are revealed. This makes it convenient to think of the measurements in a matrix form. Specifically, a non-adaptive *measurement matrix* is an $m \times n$ Boolean matrix whose i th row is the characteristic vector of the set of items participating in the i th pool, and the goal would be to design a suitable measurement matrix.

More recently, non-adaptive threshold testing has been considered by Chen and Fu [6]. They observe that a generalization of the standard notion of disjunct matrices, the latter being extensively used in the literature of classical group testing, is suitable for the threshold model. Throughout this work, we refer to this generalized notion as *strongly disjunct* matrices and to the standard notion as *classical* disjunct matrices. Using strongly disjunct matrices, they show that $O(ed^{u+1} \log(n/d))$ non-adaptive measurements suffices to identify the set of positives (within g false positives/negatives) even if up to e erroneous measurements are allowed in the model. This number of measurements almost matches (up to constant factors) the known lower bounds on the number of rows of strongly disjunct matrices. However, the dependence on the sparsity parameter is d^{u+1} , which might be prohibitive for an interesting range of parameters, when the thresholds are not too small (e.g., $\ell = u = 10$) and the sparsity parameter is rather large (e.g., $d = n^{1/10}$).

¹A vector is d -sparse if its support has size at most d .

²The original paper of Damaschke [14] uses a slightly different convention where the test returns negative if the number of positives in the group is *at most* ℓ . Accordingly he defines the gap between the thresholds to be $u - \ell - 1$, as opposed to $u - \ell$ in our paper.

In this work, we consider the non-adaptive threshold model in a possibly noisy setting, where a number of measurement outcomes (specified by an *error parameter* $e \geq 0$) may be incorrect. Our first observation is that, a new variation of classical disjunct matrices (that is in general strictly weaker than strongly disjunct matrices) suffices for the purpose of threshold group testing. Moreover, we show that this weaker notion is necessary as well, and thus, precisely captures the combinatorial structure of the non-adaptive threshold model. Using a randomness-efficient probabilistic construction (that requires $\text{poly}(d, \log n)$ bits of randomness), we construct generalized disjunct matrices with $O(d^{g+2}(\log d) \log(n/d))$ rows. Thus, we bring the exponent of d in the asymptotic number of measurements from $u + 1$ (that is optimal for strongly disjunct matrices) down to $g + 2$, which is *independent* of the actual choice of the thresholds and only depends on the *gap* between them. We also show that this tradeoff is essentially optimal.

We proceed to define a new auxiliary object, namely the notion of *regular* matrices, that turns out to be the key combinatorial object in our explicit constructions. Intuitively, given a gap $g \geq 0$, a suitable regular matrix M_1 can be used to take any measurement matrix M_2 designed for the threshold model with lower threshold $\ell = 1$ and higher threshold $u = g + 1$ and “life” it up to matrix that works for any arbitrary lower threshold $\ell' > 1$ and the same gap g . Therefore, for instance, in order to address the gap-free model, it would suffice to have a non-adaptive scheme for the classical group testing model with $\ell = u = 1$. This transformation is accomplished using a simple product that increases the height of the original matrix M_2 by a multiplicative factor equal to the height of the regular matrix M_1 , while preserving the “low-threshold” distinguishing properties of the original matrix M_2 .

Next, we introduce a framework for construction of regular matrices using *strong lossless condensers* that are fundamental objects in derandomization theory, and more generally, theoretical computer science. We show that, by using an optimal condenser, it is possible to construct regular matrices with only $O(d(\log d) \log n)$ rows. This almost matches the upper bound achieved by a probabilistic construction that we also present in this work. To this date, no explicit construction of such optimal lossless condensers is known (though probabilistic constructions are easy to come up with). However, using state of the art in explicit condensers [4, 22], we will come up with two explicit constructions of regular matrices with incomparable parameters. Namely, one with $O(d(\log d) \text{quasipoly}(\log n))$ rows and another with $O(d^{1+\beta} \text{poly}(\log n))$, where $\beta > 0$ is any arbitrary constant and the exponent of the term $\text{poly}(\log n)$ depends on the choice of β . By combining regular matrices with strongly disjunct ones (designed for the lowered thresholds $\ell' = 1$ and $u' = g + 1$), we obtain our threshold testing schemes. The bounds obtained by our final schemes are summarized in Table 1. When the lower threshold ℓ is not too small, our explicit constructions (rows M8 and M9 of Table 1) significantly improve what was previously known to be achievable even using non-constructive proofs.

The rest of the paper is organized as follows. In Section 1.1 we introduce preliminary notions and fix some notation. Section 2 reviews the notion of strongly disjunct matrices and introduces our weaker notion (for the gap-free case $g = 0$), in addition to the notion of regular matrices and its properties. In Section 3 we develop our construction of regular matrices from lossless condensers, and instantiate the parameters in Section 3.1. This in particular leads to our explicit threshold testing schemes. Finally in Section 4 we consider the case with nonzero gap, and in Section 5 we discuss the future directions.

1.1 Preliminaries

For a matrix M , we denote by $M[i, j]$ the entry of M at the i th row and the j th column. Similarly, we denote the i th entry of a vector v by $v(i)$. The *support* a vector $x \in \{0, 1\}^n$, denoted by $\text{supp}(x)$,

Table 1: Summary of the parameters achieved by various threshold testing schemes. The noise parameter $p \in [0, 1)$ is arbitrary, and thresholds $\ell, u = \ell + g$ are fixed constants. “Exp” and “Rnd” respectively indicate explicit and randomized constructions. “KS” refers to the construction of strongly disjunct matrices based on Kautz-Singleton superimposed codes [24], as described in Appendix A (the bounds in rows M1-M5 are obtained by strongly disjunct matrices).

	Number of rows	Tolerable errors	Remarks
M1	$O(d^{u+1} \frac{\log(n/d)}{(1-p)^2})$	$\Omega(pd \frac{\log(n/d)}{(1-p)^2})$	Rnd: Random strongly disjunct matrices.
M2	$O((\frac{d}{1-p})^{u+1} \log n)$	$\Omega(pd \frac{\log n}{1-p})$	Exp: KS using codes on the GV bound.
M3	$O((\frac{d \log n}{1-p})^{u+1})$	$\Omega(pd \frac{\log n}{1-p})$	Exp: KS using Reed-Solomon codes.
M4	$O((\frac{d}{1-p})^{2u+1} \log n)$	$\Omega(pd \frac{\log n}{1-p})$	Exp: KS using Algebraic Geometric codes.
M5	$O((\frac{d\sqrt{\log n}}{1-p})^{u+3/2})$	$\Omega(p(\frac{d\sqrt{\log n}}{1-p})^{3/2})$	Exp: KS using Hermitian codes ($d \gg \sqrt{\log n}$).
M6	$O(d^{g+2} \frac{(\log d) \log(n/d)}{(1-p)^2})$	$\Omega(pd \frac{\log(n/d)}{(1-p)^2})$	Rnd: Construction 2.
M7	$O(d^{g+3} \frac{(\log d) \log^2 n}{(1-p)^2})$	$\Omega(pd^2 \frac{\log^2 n}{(1-p)^2})$	Constructions 4 and 1 combined, assuming optimal condensers and strongly disjunct matrices.
M8	$O(d^{g+3} \frac{(\log d) T_2 \log n}{(1-p)^{g+2}})$	$\Omega(pd^2 \frac{T_2 \log n}{1-p})$	Exp: Constructions 4 and 1 combined using Theorem 10 and M2, where $T_2 = \exp(O(\log^3 \log n)) = \text{quasipoly}(\log n)$.
M9	$O(d^{g+3+\beta} \frac{T_3^\ell \log n}{(1-p)^{g+2}})$	$\Omega(pd^{2-\beta} \frac{\log n}{1-p})$	Exp: Constructions 4 and 1 combined using Theorem 11 and M2, where $\beta > 0$ is any arbitrary constant and $T_3 = ((\log n)(\log d))^{1+u/\beta} = \text{poly}(\log n, \log d)$.
	$\Omega(d^{g+2} \log_d n + ed^{g+1})$	e	Lower bound (see Section 4).

is a subset of $[n] := \{1, \dots, n\}$ such that $i \in \text{supp}(x)$ if and only if $x(i) = 1$. The Hamming weight of x , denote by $\text{wgt}(x)$ is defined as $|\text{supp}(x)|$. For an $m \times n$ Boolean matrix M and $S \subseteq [n]$, we denote by $M|_S$ the $m \times |S|$ submatrix of M formed by restricting M to the columns picked by S . Moreover, for a vector $x \in \{0, 1\}^n$, we use $M[x]_{\ell, u}$ to denote the set of vectors in $\{0, 1\}^m$ that correctly encode the measurement outcomes resulting from non-adaptive threshold tests defined by the measurement matrix M on x using threshold parameters ℓ, u . In the gap-free case, this set may only have a single element that we denote by $M[x]_u$. Thus, for any $y \in M[x]_{\ell, u}$ we have $y(i) = 1$ if $|\text{supp}(M_j) \cap \text{supp}(x)| \geq u$, and $y(i) = 0$ if $|\text{supp}(M_j) \cap \text{supp}(x)| < \ell$, where M_j indicates the j th row of M .

The *min-entropy* of a distribution $\mathcal{X}$ with finite support Ω is given by

$$H_\infty(\mathcal{X}) := \min_{x \in \Omega} \{-\log \mathcal{X}(x)\},$$

where $\mathcal{X}(x)$ is the probability that $\mathcal{X}$ assigns to the outcome x and logarithm is taken to base 2. A *flat* distribution is one that is uniform on its support. For such a distribution $\mathcal{X}$, we have $H_\infty(\mathcal{X}) = \log(|\text{supp}(\mathcal{X})|)$. The *statistical distance* between two distributions $\mathcal{X}$ and $\mathcal{Y}$ defined on the same finite space Ω is given by $\frac{1}{2} \sum_{s \in \Omega} |\mathcal{X}(s) - \mathcal{Y}(s)|$, which is half the ℓ_1 distance of the two distributions when regarded as vectors of probabilities over Ω . Two distributions $\mathcal{X}$ and $\mathcal{Y}$ are said to be ϵ -close if their statistical distance is at most ϵ . We will use the shorthand $\mathcal{U}_n$ for the uniform distribution on $\{0, 1\}^n$, and $X \sim \mathcal{X}$ for a random variable X drawn from a distribution $\mathcal{X}$.

The main technical tool that we use in our explicit constructions is the notion of *lossless condensers*, defined below.

Definition 1. A function $f: \{0, 1\}^{\tilde{n}} \times \{0, 1\}^t \rightarrow \{0, 1\}^{\tilde{\ell}}$ is a strong *lossless condenser* for entropy k and with *error* ϵ (in short, (k, ϵ) -condenser) if for every distribution $\mathcal{X}$ on $\{0, 1\}^{\tilde{n}}$ with min-entropy at least k , random variable $X \sim \mathcal{X}$ and a *seed* $Y \sim \mathcal{U}_t$, the distribution of $(Y, f(X, Y))$ is ϵ -close to some distribution $(\mathcal{U}_t, \mathcal{Z})$ with min-entropy at least $t + k$. A condenser is *explicit* if it is polynomial-time computable.

We will use the following “almost-injectivity” property of lossless condensers in our proofs.

Proposition 2. Let $\mathcal{X}$ be a flat distribution with min-entropy $\log K$ over a finite sample space Ω and $f: \Omega \rightarrow \Gamma$ be a mapping to a finite set Γ . If $f(\mathcal{X})$ is ϵ -close to having min-entropy $\log K$, then there is a set $T \subseteq \Gamma$ of size at least $(1 - 2\epsilon)K$ such that

$$(\forall y \in T) \quad f(x) = y \wedge f(x') = y \Rightarrow x = x'.$$

Proof. Suppose that $\mathcal{X}$ is uniformly supported on a set $S \subseteq \Omega$ of size K . For each $y \in \Gamma$, define $n_y := |\{x \in \Omega: f(x) = y\}|$. Denote by μ the distribution $f(\mathcal{X})$ over Γ and by μ' a distribution on Γ with min-entropy K that is ϵ -close to μ , which is guaranteed to exist by the assumption. Define $T := \{y \in \Gamma: n_y = 1\}$, and similarly, $T' := \{y \in \Gamma: n_y \geq 2\}$. Observe that for each $y \in \Gamma$ we have $\mu(y) = n_y/K$, and also $\text{supp}(\mu) = T \cup T'$. Thus,

$$|T| + \sum_{y \in T'} n_y = K. \tag{1}$$

The fact that μ and μ' are ϵ -close implies that

$$\sum_{y \in T'} |\mu(y) - \mu'(y)| \leq \epsilon \Rightarrow \sum_{y \in T'} (n_y - 1) \leq \epsilon K.$$

In particular, this means that $|T'| \leq \epsilon K$ (since by the choice of T' , for each $y \in T'$ we have $n_y \geq 2$). Furthermore,

$$\sum_{y \in T'} (n_y - 1) \leq \epsilon K \Rightarrow \sum_{y \in T'} n_y \leq \epsilon K + |T'| \leq 2\epsilon K.$$

This combined with (1) gives

$$|T| = K - \sum_{y \in T'} n_y \geq (1 - 2\epsilon)K$$

as desired. $\square$

2 Variations of disjunct matrices

The combinatorial structure used by Chen and Fu in their non-adaptive scheme is the following generalization of the standard notion of disjunct matrices that we refer to as *strongly disjunct* matrices throughout this work.

Definition 3. A matrix (with at least $d + u$ columns) is said to be strongly $(d, e; u)$ -disjunct if for every choice of $d + u$ columns $C_1, \dots, C_u, C'_1, \dots, C'_d$, all distinct, we have

$$|\cap_{i=1}^u \text{supp}(C_i) \setminus \cup_{i=1}^d \text{supp}(C'_i)| > e.$$

Observe that, $(d, e; u)$ -disjunct matrices are, in particular, $(d', e'; u')$ -disjunct for any $d' \leq d$, $e' \leq e$, and $u' \leq u$. Moreover, *classical* (d, e) -disjunct matrices that are extensively used in group testing literature (see [16, Ch. 7]) correspond to the special case $u = 1$.

To make the main ideas more transparent, until Section 4 we will focus on the gap-free case where $\ell = u$. The extension to nonzero gaps is straightforward and will be discussed in Section 4. Moreover, often we will implicitly assume that the Hamming weight of the Boolean vector that is to be identified is at least u (since otherwise, we know that confusions cannot be avoided), and will take the thresholds ℓ, u as fixed constants.

The notion of strongly disjunct matrices, in its general form, has been studied in the literature under different names and equivalent formulations, e.g., superimposed (u, d) -designs/codes and (u, d) cover-free families (see [5, 7, 18, 25, 34, 35] and the references therein). An important motivation for the study of this notion is the following *hidden hypergraph-learning problem* (cf. [16, Ch. 12]), itself being motivated by the so-called *complex model* in computational biology [5]: Suppose that G is a u -hypergraph³ on a vertex set V of size n , and denote by $\mathcal{V}(G)$ the set of vertices induced by the hyper-edge set of G ; i.e., $v \in \mathcal{V}(G)$ if and only if G has a hyper-edge incident to v . Then assuming that $|\mathcal{V}(G)| \leq d$ for a *sparsity parameter* d , the aim is to learn G using as few (non-adaptive) queries of the following type as possible: Each query specifies a set $Q \subseteq V$, and its corresponding answer is a Boolean value which is 1 if and only if G has a hyperedge contained in Q . It is known that [5, 20], in the hypergraph-learning problem, any suitable grouping strategy defines a strongly disjunct matrix (whose rows are characteristic vectors of individual queries Q), and conversely, any strongly disjunct matrix can be used as the incidence matrix of the set of queries. The parameter e determines “noise tolerance” of the measurement scheme. Namely, a strongly $(d, e; u)$ -disjunct matrix can uniquely distinguish between d -sparse hypergraphs even in presence of up to $\lfloor e/2 \rfloor$ erroneous query outcomes.

The key observation made by Chen and Fu [6] is that threshold group testing corresponds to the special case of the hypergraph learning problem where the hidden graph G is known to be a

³That is, a hypergraph where each edge is a set of u vertices.

u -clique⁴. In this case, the unknown Boolean vector in the corresponding threshold testing problem would be the characteristic vector of $\mathcal{V}(G)$. It follows that strongly disjunct matrices are suitable choices for the measurement matrices in threshold group testing.

Nonconstructively, a probabilistic argument akin to the standard argument for the case of classical disjunct matrices (see [16, Ch. 7]) can be used to show that strongly $(d, e; u)$ -disjunct matrices exist with $m = O(d^{u+1}(\log(n/d))/(1-p)^2)$ rows and error tolerance $e = \Omega(pd \log(n/d)/(1-p)^2)$, for any noise parameter $p \in [0, 1)$. On the negative side, however, several concrete lower bounds are known for the number of rows of such matrices [18, 34, 35]. In asymptotic terms, these results show that one must have $m = \Omega(d^{u+1} \log_d n + ed^u)$, and thus, the probabilistic upper bound is essentially optimal.

For the underlying strongly disjunct matrix, Chen and Fu [6] use a greedy construction [7] that achieves, for any $e \geq 0$, $O((e+1)d^{u+1} \log(n/d))$ rows, but may take exponential time in the size of the resulting matrix. Nevertheless, as observed by several researchers [5, 18, 20, 25], a classical explicit construction of combinatorial designs due to Kautz and Singleton [24] can be extended to construct strongly disjunct matrices. This concatenation-based construction transforms any error-correcting code having large distance into a disjunct matrix. While the original construction uses Reed-Solomon codes and achieves nice bounds, it is possible to use other families of codes. In particular, as recently shown by Porat and Rothschild [30], codes on the Gilbert-Varshamov bound (cf. [28]) result in nearly optimal disjunct matrices. Moreover, for a suitable range of parameters, they give a *deterministic* construction of such codes that runs in polynomial time in the size of the resulting disjunct matrix (albeit exponential in the dimension of the code⁵). We will elaborate on details of this class of constructions in Appendix A, and will additionally consider a family of algebraic-geometric codes and Hermitian codes which give incomparable bounds, as summarized in Table 1 (rows M2–M5).

Even though, as discussed above, the general notion of strongly $(d, e; u)$ -disjunct matrices is sufficient for threshold group testing with upper threshold u , in this section we show that a new, weaker, notion of disjunct matrices defined below (which, as we show later, turns out to be *strictly* weaker when $u > 1$), would also suffice. We also define an auxiliary notion of *regular* matrices.

Definition 4. A Boolean matrix M with n columns is called $(d, e; u)$ -regular if for every subset of columns $S \subseteq [n]$ (called the *critical set*) and every $Z \subseteq [n]$ (called the *zero set*) such that $u \leq |S| \leq d$, $|Z| \leq |S|$, $S \cap Z = \emptyset$, there are more than e rows of M at which $M|_S$ has weight exactly u and (at the same rows) $M|_Z$ has weight zero. Any such row is said to *u -satisfy* S and Z . If, in addition, for every *distinguished column* $i \in S$, more than e rows of M both *u -satisfy* S and Z and have a 1 at the i th column, the matrix is called $(d, e; u)$ -disjunct (and the corresponding “good” rows are said to *u -satisfy* i , S , and Z).

To distinguish between the above variant of disjunct matrices and strongly disjunct matrices or classical disjunct matrices, we will refer to our variant as *threshold disjunct* matrices, or (when there is no risk of confusion) simply *disjunct* matrices.

It is easy to verify that (assuming $2d \leq n$) the classical notion of $(2d-1, e)$ -disjunct matrices is equivalent to strongly $(2d-1, e; 1)$ -disjunct and $(d, e; 1)$ -disjunct. Moreover, any $(d, e; u)$ -disjunct matrix is $(d, e; u)$ -regular, $(d-1, e; u-1)$ -regular, and classical (d, e) -disjunct (but the reverse implications do not in general hold). Therefore, the above-mentioned lower bound of

⁴A u -clique on the vertex set V is a u -hypergraph (V, E) such that, for some $V' \subseteq V$, E is the set of all subsets of V' of size u .

⁵In this regard, this construction of disjunct matrices can be considered *weakly explicit* in that, contrary to fully explicit constructions, it is not clear if each individual entry of the matrix can be computed in time $\text{poly}(d, \log n)$.

$m = \Omega(d^2 \log_d n + ed)$ that applies for (d, e) -disjunct matrices holds for $(d, e; u)$ -disjunct matrices as well. Below we show that our notion of disjunct matrices is necessary and sufficient for the purpose of threshold group testing:

Lemma 5. *Let M be an $m \times n$ Boolean matrix that is $(d, e; u)$ -disjunct. Then for every distinct d -sparse vectors $x, x' \in \{0, 1\}^n$ such that⁶ $\text{supp}(x) \not\subseteq \text{supp}(x')$, $\text{wgt}(x) \geq |\text{supp}(x') \setminus \text{supp}(x)|$ and $\text{wgt}(x) \geq u$, we have*

$$|\text{supp}(M[x]_u) \setminus \text{supp}(M[x']_u)| > e. \quad (2)$$

Conversely, if M satisfies (2) for every choice of x and x' as above, it must be $(\lfloor d/2 \rfloor, e; u)$ -disjunct.

Proof. First, suppose that M is $(d, e; u)$ -disjunct, and let $y := M[x]_u$ and $y' := M[x']_u$. Take any $i \in \text{supp}(x) \setminus \text{supp}(x')$, and let $S := \text{supp}(x)$ and $Z := \text{supp}(x') \setminus \text{supp}(x)$. Note that $|S| \leq d$ and by assumption, we have $|Z| \leq |S|$. Now, Definition 4 implies that there is a set E of more than e rows of M that u -satisfy i as the distinguished column, S as the critical set and Z as the zero set. Thus for every $j \in E$, the j th row of M restricted to the columns chosen by $\text{supp}(x)$ must have weight exactly u , while its weight on $\text{supp}(x')$ is less than u . Therefore, $y(j) = 1$ and $y'(j) = 0$ for more than e choices of j .

For the converse, consider any choice of a distinguished column $i \in [n]$, a critical set $S \subseteq [n]$ containing i (such that $|S| \geq u$), and a zero set $Z \subseteq [n]$ where $|Z| \leq |S|$. Define d -sparse Boolean vectors $x, x' \in \{0, 1\}^n$ so that $\text{supp}(x) := S$ and $\text{supp}(x') := Z \cup (S \setminus \{i\})$. Let $y := M[x]_u$ and $y' := M[x']_u$ and $E := \text{supp}(y) \setminus \text{supp}(y')$. By assumption we know that $|E| > e$. Take any $j \in E$. Since $y(j) = 1$ and $y'(j) = 0$, we get that the j th row of M restricted to the columns picked by $Z \cup (S \setminus \{i\})$ must have weight at most $u - 1$, whereas it must have weight at least u when restricted to S . As the sets $\{i\}$, $S \setminus \{i\}$, and Z are disjoint, this can hold only if $M[j, i] = 1$, and moreover, the j th row of M restricted to the columns picked by S (resp., Z) has weight exactly u (resp., zero). Hence, this row (as well as all the rows of M picked by E) must u -satisfy i , S , and Z , confirming that M is $(\lfloor d/2 \rfloor, e; u)$ -disjunct. $\square$

We will use regular matrices as intermediate building blocks in our constructions of disjunct matrices to follow. The connection with disjunct matrices is made apparent through a direct product of matrices defined in Construction 1. Intuitively, using this product, regular matrices can be used to transform any measurement matrix suitable for the standard group testing model to one with comparable properties in the threshold model. The following lemma formalizes the idea.

Lemma 6. *Let M_1 and M_2 be Boolean matrices with n columns, such that M_1 is $(d - 1, e_1; u - 1)$ -regular. Let $M := M_1 \odot M_2$, and suppose that for d -sparse Boolean vectors $x, x' \in \{0, 1\}^n$ such that*

⁶ Observe that at least one of the two possible orderings of any two distinct d -sparse vectors, at least one having weight u or more, satisfies this condition.

- *Given:* Boolean matrices M_1 and M_2 that are $m_1 \times n$ and $m_2 \times n$, respectively.
- *Output:* An $m \times n$ Boolean matrix $M_1 \odot M_2$, where $m := m_1 m_2$.
- *Construction:* Let the rows of $M := M_1 \odot M_2$ be indexed by the set $[m_1] \times [m_2]$. Then the row corresponding to (i, j) is defined as the bit-wise or of the i th row of M_1 and the j th row of M_2 .

Construction 1: Direct product of measurement matrices.

$\text{wgt}(x) \geq \text{wgt}(x')$, we have

$$|\text{supp}(M_2[x]_1) \setminus \text{supp}(M_2[x']_1)| \geq e_2.$$

Then, $|\text{supp}(M[x]_u) \setminus \text{supp}(M[x']_u)| \geq (e_1 + 1)e_2$.

Proof. First we consider the case where $u > 1$. Let $y := M_2[x]_1 \in \{0, 1\}^{m_2}$, $y' := M_2[x']_1 \in \{0, 1\}^{m_2}$, where m_2 is the number of rows of M_2 , and let $E := \text{supp}(y) \setminus \text{supp}(y')$. By assumption, $|E| \geq e_2$. Fix any $i \in E$ so that $y(i) = 1$ and $y'(i) = 0$. Therefore, the i th row of M_2 must have all zeros at positions corresponding to $\text{supp}(x')$ and there is a $j \in \text{supp}(x) \setminus \text{supp}(x')$ such that $M_2[i, j] = 1$. Define $S := \text{supp}(x) \setminus \{j\}$, $Z := \text{supp}(x') \setminus \text{supp}(x)$, $z := M[x]_u$ and $z' := M[x']_u$.

As $\text{wgt}(x) \geq \text{wgt}(x')$, we know that $|Z| \leq |S| + 1$. The extreme case $|Z| = |S| + 1$ only happens when x and x' have disjoint supports, in which case one can remove an arbitrary element of Z to ensure that $|Z| \leq |S|$ and the following argument (considering the assumption $u > 1$) still goes through.

By the definition of regularity, there is a set E_1 consisting of at least $e_1 + 1$ rows of M_1 that $(u - 1)$ -satisfy the critical set S and the zero set Z . Pick any $k \in E_1$, and observe that z must have a 1 at position (k, i) . This is because the row of M indexed by (k, i) has a 1 at the j th position (since the k th row of M_2 does), and at least $u - 1$ more 1's at positions corresponding to $\text{supp}(x) \setminus \{j\}$ (due to regularity of M_1). On the other hand, note that the k th row of M_1 has at most $u - 1$ ones at positions corresponding to $\text{supp}(x')$ (because $\text{supp}(x') \subseteq S \cup Z$), and the i th row of M_2 has all zeros at those positions (because $y'(i) = 0$). This means that the row of M indexed by (k, i) (which is the bit-wise or of the k th row of M_1 and the i th row of M_2) must have less than u ones at positions corresponding to $\text{supp}(x')$, and thus, z' must be 0 at position (k, i) . Therefore, z and z' differ at position (k, i) .

Since there are at least e_2 choices for i , and for each choice of i , at least $e_1 + 1$ choices for k , we conclude that in at least $(e_1 + 1)e_2$ positions, z has a one while z' has a zero.

The argument for $u = 1$ is similar, in which case it suffices to take $S := \text{supp}(x)$ and $Z := \text{supp}(x') \setminus \text{supp}(x)$. $\square$

As a corollary it follows that, when M_1 is a $(d - 1, e_1; u - 1)$ -regular and M_2 is a (d, e_2) -disjunct matrix, the product $M := M_1 \odot M_2$ will distinguish between any two distinct d -sparse vectors (of weight at least u) in at least $(e_1 + 1)(e_2 + 1)$ positions of the measurement outcomes. This combined with Lemma 5 would imply that M is, in particular, $(\lfloor d/2 \rfloor, (e_1 + 1)(e_2 + 1) - 1; u)$ -disjunct. However, using a direct argument similar to the above lemma it is possible to obtain a slightly better result, given by Lemma 7.

Lemma 7. *Suppose that M_1 is a $(d, e_1; u - 1)$ -regular and M_2 is a $(2d, e_2)$ -disjunct matrix. Then $M_1 \odot M_2$ is a $(d, (e_1 + 1)(e_2 + 1) - 1; u)$ -disjunct matrix.* $\square$

As a particular example of where Lemma 6 can be used, we remark that the measurement matrices constructed in [9] that are not necessarily disjunct but allow approximation of sparse vectors in highly noisy settings of the standard group testing model (as well as those used in adaptive two-stage schemes; cf. [8] and the references therein), can be combined with regular matrices to offer the same qualities in the threshold model. In the same way, numerous existing results in group testing can be ported to the threshold model by using Lemma 6.

3 Constructions

In this section, we obtain several construction of regular and disjunct matrices. Our first construction, described in Construction 2, is a randomness-efficient probabilistic construction that can be

analyzed using standard techniques from the probabilistic method. The bounds obtained by this construction are given in Lemma 8 below. The amount of random bits required by this construction is polynomially bounded in d and $\log n$, which is significantly smaller than it would be had we picked the entries of M fully independently.

Lemma 8. *For every $p \in [0, 1)$ and integer parameter $u > 0$, Construction 2 with $m' = O_u(d \log(n/d)/(1-p)^2)$ (resp., $m' = O_u(d^2 \log(n/d)/(1-p)^2)$) outputs a $(d, \Omega_u(pm'); u)$ -regular (resp., $(d, \Omega_u(pm'/d); u)$ -disjunct) matrix with probability $1 - o(1)$.*

Proof. We show the claim for regular matrices, the proof for disjunct matrices is similar. Consider any particular choice of a critical set $S \subseteq [n]$ and a zero set $Z \subseteq [n]$ such that $u \leq |S| \leq d$ and $|Z| \leq |S|$. Choose an integer i so that $2^{i-1}u \leq |S| \leq 2^i u$, and take any $j \in [m']$. Denote the (i, j) th row of M by the random variable $\mathbf{w} \in \{0, 1\}^n$, and by q the “success” probability that $\mathbf{w}|_S$ has weight exactly u and $\mathbf{w}|_Z$ is all zeros. For an integer $\ell > 0$, we will use the shorthand 1^ℓ (resp., 0^ℓ) for the all-ones (resp., all-zeros) vector of length ℓ . We have

$$\begin{aligned}
q &= \sum_{\substack{R \subseteq [S] \\ |R|=u}} \Pr[(\mathbf{w}|_R) = 1^u \wedge (\mathbf{w}|_{Z \cup (S \setminus R)}) = 0^{|S|+|Z|-u}] \\
&= \sum_R \Pr[(\mathbf{w}|_R) = 1^u] \cdot \Pr[(\mathbf{w}|_{Z \cup (S \setminus R)}) = 0^{|S|+|Z|-u} \mid (\mathbf{w}|_R) = 1^u] \\
&\stackrel{(a)}{=} \sum_R (1/(2^{i+2}u))^u \cdot (1 - \Pr[(\mathbf{w}|_{Z \cup (S \setminus R)}) \neq 0^{|S|+|Z|-u} \mid (\mathbf{w}|_R) = 1^u]) \\
&\stackrel{(b)}{\geq} \sum_R (1/(2^{i+2}u))^u \cdot (1 - (|S| + |Z| - u)/(2^{i+2}u)) \\
&\geq \frac{1}{2} \binom{|S|}{u} (1/(2^{i+2}u))^u \geq \frac{1}{2} \left(\frac{|S|}{u} \right)^u \cdot (1/(2^{i+2}u))^u \geq \frac{1}{2^{3u+1} \cdot u^u} =: c, \tag{3}
\end{aligned}$$

where (a) and (b) use the fact that the entries of $\mathbf{w}$ are $(u+1)$ -wise independent, and (b) uses an additional union bound. Here the lower bound $c > 0$ is a constant that only depends on u . Now, let $e := m'pq$. using Chernoff bounds, and independence of the rows, the probability that there are at most e rows (among $(i, 1), \dots, (i, m')$) whose restrictions to S and Z have weights u and 0, respectively, becomes upper bounded by

$$\exp(-(m'q - e)^2/(2m'q)) = \exp(-(1-p)^2 m'q/2) \leq \exp(-(1-p)^2 m'c/2).$$

Now take a union bound on all the choices of S and Z to conclude that the probability that the

- *Given:* Integer parameters n, m', d, u .
- *Output:* An $m \times n$ Boolean matrix M , where $m := m' \lceil \log(d/u) \rceil$.
- *Construction:* Let $r := \lceil \log(d/u) \rceil$. Index the rows of M by $[r] \times [m']$. Sample the (i, j) th row of M independently from a $(u+1)$ -wise independent distribution on n bit vectors, where each individual bit has probability $1/(2^{i+2}u)$ of being 1.

Construction 2: Probabilistic construction of regular and disjunct matrices.

resulting matrix is not $(d, e; u)$ -regular is at most

$$\left(\sum_{s=u}^d \binom{n}{s} \sum_{z=0}^s \binom{n-s}{z} \right) \exp(-(1-p)^2 m' c/2),$$

which can be made $o(1)$ by choosing $m' = O_u(d \log(n/d)/(1-p)^2)$.

The proof of the claim for disjunct matrices follows along the same lines, except that we additionally need the vector $\mathbf{w}$ to be 1 at the position corresponding to the distinguished column i . Under this additional requirement, the lower bound on q would become $\Omega_u(1/d)$, and this only increases the number of rows by a factor $O_u(d)$. $\square$

A significant part of this work is a construction of regular matrices using strong lossless condensers. Details of the construction are described in Construction 4 that assumes a family of lossless condensers with different entropy requirements⁷, and in turn, uses Construction 3 as a building block. The theorem below analyzes the obtained parameters without specifying any particular choice for the underlying family of condensers.

- *Given:* A strong lossless (k, ϵ) -conductor $f: \{0, 1\}^{\tilde{n}} \times \{0, 1\}^t \rightarrow \{0, 1\}^{\tilde{\ell}}$, integer parameter $u \geq 1$ and real parameter $p \in [0, 1)$ such that $\epsilon < (1-p)/16$,
- *Output:* An $m \times n$ Boolean matrix M , where $n := 2^{\tilde{n}}$ and $m = 2^{t+k} O_u(2^{u(\tilde{\ell}-k)})$.
- *Construction:* Let $G_1 = (\{0, 1\}^{\tilde{\ell}}, \{0, 1\}^k, E_1)$ be any bipartite bi-regular graph with left vertex set $\{0, 1\}^{\tilde{\ell}}$, right vertex set $\{0, 1\}^k$, edge set E_1 , left degree $d_\ell := 8u$, and right degree $d_r := 8u2^{\tilde{\ell}-k}$. Replace each right vertex v of G_1 with $\binom{d_r}{u}$ vertices, one for each subset of size u of the vertices on the neighborhood of v , and connect them to the vertices in the corresponding subsets. Denote the resulting graph by $G_2 = (\{0, 1\}^{\tilde{\ell}}, V_2, E_2)$, where $|V_2| = 2^k \binom{d_r}{u}$ and E_2 is the edge set of the graph. Define the bipartite graph $G_3 = (\{0, 1\}^{\tilde{n}}, V_3, E_3)$, where $V_3 := \{0, 1\}^t \times V_2$ is the set of right vertices, as follows: Each left vertex $x \in \{0, 1\}^{\tilde{n}}$ is connected to $(y, \Gamma_2(f(x, y)))$, for each $y \in \{0, 1\}^t$, where $\Gamma_2(\cdot)$ denotes the neighborhood function of G_2 (i.e., $\Gamma_2(v)$ denotes the set of vertices adjacent to v in G_2). The output matrix M is the bipartite adjacency matrix of G_3 with columns indexed by the left vertices of row indexed by the right vertices of the graph.

Construction 3: A building block for construction of regular matrices.

Theorem 9. *The $m \times n$ matrix M output by Construction 4 is $(d, p\gamma 2^t; u)$ -regular, where $\gamma = \max\{1, \Omega_u(d \cdot \min\{2^{k(i)-\tilde{\ell}(i)} : i = 0, \dots, r\})\}$.*

Proof. As a first step, we verify the upper bound on the number of measurements m . Each matrix M_i has $m_i = 2^{t+k(i)} O_u(2^{u(\tilde{\ell}(i)-k(i))})$ rows, and M'_i has $m_i r_i$ rows, where $r_i = 2^{r-i}$. Therefore, the number of rows of M is

$$\sum_{i=0}^r r_i m_i = \sum_{i=0}^r 2^{t+\log u' + r+1} m_i = 2^t d \sum_{i=0}^r O_u(2^{u(\tilde{\ell}(i)-k(i))}).$$

⁷We have assumed that all the functions in the family have the same seed length t . If this is not the case, one can trivially set t to be the largest seed length in the family.

- *Given:* Integer parameters $d \geq u \geq 1$, real parameter $p \in [0, 1)$, and a family $f_0, \dots, f_r$ of strong lossless condensers, where $r := \lceil \log(d/u') \rceil$ and u' is the smallest power of two such that $u' \geq u$. Each $f_i: \{0, 1\}^{\tilde{n}} \times \{0, 1\}^t \rightarrow \{0, 1\}^{\tilde{\ell}(i)}$ is assumed to be a strong lossless $(k(i), \epsilon)$ -condenser, where $k(i) := \log u' + i + 1$ and $\epsilon < (1 - p)/16$.
- *Output:* An $m \times n$ Boolean matrix M , where $n := 2^{\tilde{n}}$ and $m = 2^t d \sum_{i=0}^r O_u(2^{u(\tilde{\ell}(i) - k(i))})$.
- *Construction:* For each $i \in \{0, \dots, r\}$, denote by M_i the output matrix of Construction 3 when instantiated with f_i as the underlying condenser, and by m_i its number of rows. Define $r_i := 2^{r-i}$ and let M'_i denote the matrix obtained from M_i by repeating each row r_i times. Construct the output matrix M by stacking $M'_0, \dots, M'_r$ on top of one another.

Construction 4: Regular matrices from strong lossless condensers.

Let $S, Z \subseteq \{0, 1\}^{\tilde{n}}$ respectively denote any choice of a critical set and zero set of size at most d , where $|Z| \leq |S|$, and choose an integer $i \geq 0$ so that $2^{i-1}u' \leq |S| \leq 2^i u'$. Arbitrarily grow the two sets S and Z to possibly larger, and disjoint, sets $S' \supseteq S$ and $Z' \supseteq Z$ such that $|S'| = |Z'| = 2^i u'$ (for simplicity we have assumed that $d \leq n/2$). Our goal is to show that there are “many” rows of the matrix M_i (in Construction 4) that u -satisfy S and Z .

Let $k := k(i) = \log u' + i + 1$, $\tilde{\ell} := \tilde{\ell}(i)$, and denote by G_1, G_2, G_3 the bipartite graphs used by the instantiation of Construction 3 that outputs M_i . Thus we need to show that “many” right vertices of G_3 are each connected to exactly u of the vertices in S and none of those in Z .

Consider the uniform distribution $\mathcal{X}$ on the set $S' \cup Z'$, which has min-entropy $\log u' + i + 1$. By an averaging argument, since the condenser f_i is strong, for more than a p fraction of the choices of the seed $y \in \{0, 1\}^t$ (call them *good seeds*), the distribution $\mathcal{Z}_y := f_i(\mathcal{X}, y)$ is $\epsilon/(1 - p)$ -close (in particular, $1/16$ -close) to a distribution with min-entropy $\log u' + i + 1$.

Fix any good seed $y \in \{0, 1\}^t$. Let $G = (\{0, 1\}^{\tilde{n}}, \{0, 1\}^{\tilde{\ell}}, E)$ denote a bipartite graph representation of f_i , where each left vertex $x \in \{0, 1\}^{\tilde{n}}$ is connected to $f_i(x, y)$ on the right. Denote by $\Gamma_y(S' \cup Z')$ the right vertices of G corresponding to the neighborhood of the set of left vertices picked by $S' \cup Z'$. Note that $\Gamma_y(S' \cup Z') = \text{supp}(\mathcal{Z}_y)$. Using Proposition 2, we see that since $\mathcal{Z}_y$ is $1/16$ -close to having min-entropy $\log(|S' \cup Z'|)$, there are at least $(7/8)|S' \cup Z'|$ vertices in $\Gamma(S' \cup Z')$ that are each connected to exactly one left vertex in $S' \cup Z'$. Since $|S| \geq |S' \cup Z'|/4$, this implies that at least $|S' \cup Z'|/8$ vertices in $\Gamma(S' \cup Z')$ (call them Γ'_y) are connected to exactly one left vertex in S and no other vertex in $S' \cup Z'$. In particular we get that $|\Gamma'_y| \geq 2^{k-3}$.

Now, in G_1 , let T_y be the set of left vertices corresponding to Γ'_y (regarding the left vertices of G_1 in one-to-one correspondence with the right vertices of G). The number of edges going out of T_y in G_1 is $d_\ell |T_y| \geq u 2^k$. Therefore, as the number of the right vertices of G_1 is 2^k , there must be at least one right vertex that is connected to at least u vertices in T_y . Moreover, a counting argument shows that the number of right vertices connected to u or more vertices in T_y is at least $2^{k-\tilde{\ell}} 2^k / (10u)$.

Observe that in construction of G_2 from G_1 , any right vertex of G_1 is replicated $\binom{d_r}{u}$ times, one for each u -subset of its neighbors. Therefore, for a right vertex of G_1 that is connected to *at least* u left vertices in T_y , one or more of its copies in G_2 must be connected to *exactly* u vertex in T_y (among the left vertices of G_2) and no other vertex (since the right degree of G_2 is equal to u).

Define $\gamma' := \max\{1, 2^{k-\tilde{\ell}} 2^k / (10u)\}$. From the previous argument we know that, looking at T_y as a set of left vertices of G_2 , there are at least γ' right vertices on the neighborhood of T_y in G_2 that are connected to exactly u of the vertices in T_y and none of the left vertices outside T_y . Letting

v_y be any such vertex, this implies that the vertex $(y, v_y) \in V_3$ on the right part of G_3 is connected to exactly u of the vertices in S , and none of the vertices in Z . Since the argument holds for every good seed y , the number of such vertices is at least the number of good seeds, which is more than $p\gamma'2^t$. Since the rows of the matrix m_i are repeated $r_i = 2^{r-i}$ times in M , we conclude that M has at least $p\gamma'2^{t+r-i} \geq p\gamma'2^t$ rows that u -satisfy S and Z , and the claim follows. $\square$

3.1 Instantiations

We now instantiate the result obtained in Theorem 9 by various choices of the family of lossless condensers. The crucial factors that influence the number of measurements are the seed length and the output length of the condenser.

Non-constructively, it can be shown that strong (k, ϵ) lossless condensers with input length $\tilde{n}$, seed length $t = \log \tilde{n} + \log(1/\epsilon) + O(1)$, and output length $\tilde{\ell} = k + \log(1/\epsilon) + O(1)$ exist, and moreover, almost matching lower bounds are known [4]. In fact, the optimal parameters can be achieved by a random function with overwhelming probability. In this work, we consider two important explicit constructions of lossless condensers. Namely, one based on “zig-zag products” due to Capalbo et al. [4] and another, coding theoretic, construction due to Guruswami et al. [22].

Theorem 10. [4] *For every $k \leq \tilde{n} \in \mathbb{N}$, $\epsilon > 0$ there is an explicit lossless (k, ϵ) condenser with seed length $O(\log^3(\tilde{n}/\epsilon))$ and output length $k + \log(1/\epsilon) + O(1)$.*

Theorem 11. [22] *For all constants $\alpha \in (0, 1)$ and every $k \leq \tilde{n} \in \mathbb{N}$, $\epsilon > 0$ there is an explicit strong lossless (k, ϵ) condenser with seed length $t = (1 + 1/\alpha) \log(\tilde{n}k/\epsilon) + O(1)$ and output length $\tilde{\ell} = t + (1 + \alpha)k$.*

As a result, we use Theorem 9 with the above condensers to obtain the following.

Theorem 12. *Let $u > 0$ be fixed, and $p \in [0, 1)$ be a real parameter. Then for integer parameters $d, n \in \mathbb{N}$ where $u \leq d \leq n$,*

1. *Using an optimal lossless condenser in Construction 4 results in an $m_1 \times n$ matrix M_1 that is $(d, e_1; u)$ -regular, where $m_1 = O(d(\log n)(\log d)/(1-p)^{u+1})$ and $e_1 = \Omega(pd \log n)$,*
2. *Using the lossless condenser of Theorem 10 in Construction 4 results in an $m_2 \times n$ matrix M_2 that is $(d, e_2; u)$ -regular, where $m_2 = O(T_2 d(\log d)/(1-p)^u)$ for some $T_2 = \exp(O(\log^3((\log n)/(1-p)))) = \text{quasipoly}(\log n)$, and $e_2 = \Omega(pd T_2(1-p))$.*
3. *Let $\beta > 0$ be any fixed constant. Then Construction 4 can be instantiated using the lossless condenser of Theorem 11 so that we obtain an $m_3 \times n$ matrix M_3 that is $(d, e_3; u)$ -regular, where $m_3 = O(T_3^{1+u} d^{1+\beta} (\log d))$ for $T_3 := ((\log n)(\log d)/(1-p))^{1+u/\beta} = \text{poly}(\log n, \log d)$, and $e_3 = \Omega(p \max\{T_3, d^{1-\beta/u}\})$.*

Proof. First we show the claim for M_1 . In this case, we take each f_i in Construction 4 to be an optimal lossless condenser satisfying the (non-constructive) bounds obtained in⁸ [4]. Thus we have that $2^t = O(\tilde{n}/\epsilon) = O(\log n/\epsilon)$, and for every $i = 0, \dots, r$, we have $2^{\tilde{\ell}(i)-k(i)} = O(1/\epsilon)$, where $\epsilon = O(1-p)$. Now we apply Theorem 9 to obtain the desired bounds (and in particular, $\gamma = \Omega(\epsilon d)$).

Similarly, for the construction of M_2 we set up each f_i with the explicit construction of condensers in Theorem 10 for min-entropy $k(i)$. In this case, the maximum required seed length

⁸This result is similar in spirit to the probabilistic argument used in [31] for showing the existence of good extractors.

is $t = O(\log^3(\tilde{n}/\epsilon))$, and we let $T_2 := 2^t = \exp(O(\log^3((\log n)/(1-p))))$. Moreover, for every $i = 0, \dots, r$, we have $2^{\tilde{\ell}(i)-k(i)} = O(1/\epsilon)$. Plugging these parameters in Theorem 9 gives $\gamma = \Omega(\epsilon d)$ and the bounds on m_2 and e_2 follow.

Finally, for M_3 we use Theorem 11 with $\alpha := \beta/u$. Thus the maximum seed length becomes $t = (1 + u/\beta) \log(\tilde{n}(\log d)/(1-p)) + O(1)$, and for every $i = 0, \dots, r$, we have $\tilde{\ell}(i) - k(i) = O(t + \beta(\log d)/u)$. Clearly, $T_3 = \Theta(2^t)$, and thus (using Theorem 9) the number of measurements becomes $m_3 = T^{1+u} d^{1+\beta} (\log d)$. Moreover, we get $\gamma = \max\{1, \Omega(d^{1-\beta/u}/T)\}$, which gives $e_3 = \Omega(pT\gamma) = p \max\{T, d^{1-\beta/u}\}$, as claimed. $\square$

By combining this result with Lemma 7 using any explicit construction of classical disjunct matrices, we obtain $(d, e; u)$ -disjunct matrices that can be used in the threshold model with any fixed threshold, sparsity d , and error tolerance $\lfloor e/2 \rfloor$. In particular, using the coding-theoretic explicit construction of nearly optimal classical disjunct matrices from codes on the Gilbert-Varshamov bound [30] (Theorem 19 in the appendix), we obtain $(d, e; u)$ -disjunct matrices with $m = O(m' d^2 (\log n)/(1-p)^2)$ rows and error tolerance $e = \Omega(e' p d (\log n)/(1-p))$, where m' and e' are respectively the number of rows and error tolerance of any of the regular matrices obtained in Theorem 12. We note that in all cases, the final dependence on the sparsity parameter d is, roughly, $O(d^3)$ which has an exponent independent of the threshold u . Rows M7–M9 of Table 1 summarize the obtained parameters for the general case (with arbitrary gaps). We see that, when d is not negligibly small (e.g., $d = n^{1/10}$), the bounds obtained by our explicit constructions are significantly better than those offered by strongly disjunct matrices.

4 The case with positive gaps

In preceding sections we have on the case where $g = 0$. However, in this section we observe that all the techniques that we have developed in this work can be extended to the positive-gap case in a straightforward way. The main observations are listed below. Recall from [14] that in the positive-gap case, we can only hope to distinguish between distinct d -sparse vectors x and x' where at least one has support size u or more and either $|\text{supp}(x) \setminus \text{supp}(x')| > g$ or $|\text{supp}(x') \setminus \text{supp}(x)| > g$. We will call any pair of such vectors *distinguishable*.

Observation 1

For the positive-gap case, Definition 4 of threshold disjunct matrices can be adapted to allow more than one distinguished column in disjunct matrices. In particular, in general we may require the matrix M to have more than e rows that u -satisfy every choice of a critical set S , a zero set Z , and any set of $g+1$ designated columns $I \subseteq S$ (at which all entries of the corresponding rows must be 1). Denote this generalized notion by $(d, e; u, g)$ -disjunct matrices. It is straightforward to extend the arguments of Lemma 5 to show that the generalized notion of $(d, e; u, g)$ -disjunct matrices is necessary and sufficient to capture non-adaptive threshold group testing with upper threshold u and gap g . More precisely, the generalized definitions of threshold disjunct and regular matrices are as follows.

Definition 13 (Definition 4, generalized). Let n, d, e, u, g be non-negative integers where $g < u \leq d \leq n$. A Boolean matrix M with n columns is called $(d, e; u, g)$ -disjunct if for every subset of columns $S \subseteq [n]$ (called the *critical set*), every $Z \subseteq [n]$ (called the *zero set*) such that $u \leq |S| \leq d$, $|Z| \leq |S|$, $S \cap Z = \emptyset$, and every set $I \subseteq S$ of $g+1$ distinguished columns ($|I| = g+1$), there are more than e rows of M that u -satisfy S and Z and moreover, $M|_I$ has all ones at those columns.

Moreover, M is called $(d, e; u, g)$ -regular if for every choice of the critical and zero sets $S, Z \subseteq [n]$ with $|Z| \leq |S| + g$, there is a set of more than e rows of M that $(u - g)$ -satisfy S and Z .

Note the slight difference between the notion of regular matrices above compared to Definition 4, namely, that the zero set Z can now be slightly larger than the critical set S (by at most u), and that the matrix is now required to $(u - g)$ -satisfy (as opposed to u -satisfy) every choice of S and Z . The two notions coincide for $g = 0$. In general, the difference between the two notions of regular matrices is negligible as long as the parameter g remains small. In particular, it is straightforward to verify that all our results about the construction of regular matrices in the gap-free case (Constructions 2 and 4) as well as the obtained bounds (Lemma 8, Theorem 9 and Theorem 12) hold for the generalized notion of regular matrices with only a slight effect on the hidden terms that only depend on the threshold parameter u . We will see, however, that the generalized notion of threshold-disjunct matrices is stronger than Definition 4 and the extra requirements may substantially affect the bounds (but not the construction techniques).

Below we show that the generalized notion of threshold disjunct matrices precisely captures the combinatorial structure needed for threshold group testing with arbitrary gap.

Lemma 14 (Lemma 5, generalized). *Let M be an $m \times n$ Boolean matrix that is $(d, e; u, g)$ -disjunct, and define $\ell := u - g$. Then for every distinguishable d -sparse vectors $x, x' \in \{0, 1\}^n$, each having support size u or more and such that⁹ $|\text{supp}(x) \setminus \text{supp}(x')| > g$ and $\text{wgt}(x) \geq |\text{supp}(x') \setminus \text{supp}(x)|$, the following holds. Let $y \in M[x]_{\ell, u}$ and $y' \in M[x']_{\ell, u}$. Then,*

$$|\text{supp}(y) \setminus \text{supp}(y')| > e. \quad (4)$$

Conversely, if M satisfies (4) for every choice of x, x', y, y' as above, it must be $(\lfloor d/2 \rfloor, e; u, g)$ -disjunct (assuming $n > d + g$).

Proof. First, suppose that M is $(d, e; u, g)$ -disjunct, and let $y := M[x]_u$ and $y' := M[x']_u$. Take any $I \subseteq \text{supp}(x) \setminus \text{supp}(x')$ of size $g + 1$, and let $S := \text{supp}(x)$ and $Z := \text{supp}(x') \setminus \text{supp}(x)$. Note that $|S| \leq d$ and by assumption, we have $|Z| \leq |S|$. Now, Definition 13 implies that there is a set E of more than e rows of M that u -satisfy I as the set of distinguished columns, S as the critical set and Z as the zero set. Thus for every $j \in E$, the j th row of M restricted to the columns chosen by $\text{supp}(x)$ must have weight exactly u , while its weight on $\text{supp}(x')$ is less than $u - g = \ell$. Therefore, $y(j) = 1$ and $y'(j) = 0$ for more than e choices of j .

For the converse, consider any choice of a set of distinguished columns $I \subseteq [n]$ with $|I| = g + 1$, a critical set $S \subseteq [n]$ containing I (such that $|S| \geq u$), and a zero set $Z \subseteq [n]$ where $|Z| \leq |S|$. Define d -sparse Boolean vectors $x, x' \in \{0, 1\}^n$ so that $\text{supp}(x) := S$ and $\text{supp}(x') := Z \cup (S \setminus I)$. We note that $|\text{supp}(x)| \geq u$ and also, without loss of generality, $|\text{supp}(x')| \geq u$ (if the latter is not the case, we can simply enlarge Z by arbitrarily adding up to $g + 1$ elements outside the support of S to it and observe that it suffices to show the claim for the larger Z).

Let $y \in M[x]_{\ell, u}$ and $y' \in M[x']_{\ell, u}$ and $E := \text{supp}(y) \setminus \text{supp}(y')$. By assumption we know that $|E| > e$. Take any $j \in E$. Since $y(j) = 1$ and $y'(j) = 0$, we get that the j th row of M restricted to the columns picked by $Z \cup (S \setminus I)$ must have weight at most $\ell - 1 = u - (g + 1)$, whereas it must have weight at least u when restricted to S . As the sets $I, S \setminus I$, and Z are disjoint and $|I| = g + 1$, this can hold only if the j th row of M restricted to the columns picked by S, Z , and I has weights exactly $u, 0$, and $g + 1$, respectively. Hence, this row (as well as all the rows of M picked by E) must u -satisfy I, S , and Z , confirming that M is $(\lfloor d/2 \rfloor, e; u)$ -disjunct. $\square$

⁹ As in Lemma 5, one can verify that at least one of the two possible orderings of any pair of distinguishable vectors satisfies this condition.

Note that, for technical reasons, the above lemma assumes both vectors x and x' to have support size u or more. This is no loss of generality since, if $|\text{supp}(x)| \geq u$ while $|\text{supp}(x')| < u$, we have $M_{\ell,u}[x'] = 0$ and one can trivially use the disjunctness property of M and show that $|\text{supp}(M_{\ell,u}[x])| > e$.

Observation 2

The following proposition directly follows from the definitions, and relates strongly disjunct matrices to generalized threshold disjunct matrices.

Proposition 15. *Let n, d, e, u, g be non-negative integers where $g < u \leq d \leq n - (d + g + 1)$. Suppose that M and M' are binary $m \times n$ matrices, where M is $(d, e; u, g)$ -disjunct and M' is strongly $(2d, e; u)$ -disjunct. Then, M is strongly $(d, e; g + 1)$ -disjunct and M' is $(d, e; u, g)$ -disjunct.*

Proof. First, we verify the conditions of Definition 3 for M . Consider any pair of disjoint sets $I, Z \subseteq [n]$ where $|I| = g + 1$ and $|Z| \leq d$. Let $S \subseteq [n]$ be any set of size d containing I and disjoint from Z . Note that $|Z| \leq |S|$. From Definition 13 (with the critical set S , zero set Z , and distinguished set I), there is a set of more than e rows of M at which $M|_Z$ is all zeros and $M|_I$ is all ones. In other words, denoting the i th column of M by C_i , we have that

$$|\cap_{i \in I} \text{supp}(C_i) \setminus \cup_{i \in Z} \text{supp}(C_i)| > e,$$

as required by Definition 3.

Now consider the matrix M' and any choice of a I, S, Z as in Definition 13. Let $J \subseteq S$ be any subset of S of size u that contains I , and $S' := Z \cup (S \setminus J)$. Note that $|S'| \leq |S| + |Z| \leq 2d$. Now from Definition 3 of strongly disjunct matrices, we know that

$$|\cap_{i \in J} \text{supp}(C_i) \setminus \cup_{i \in S'} \text{supp}(C_i)| > e.$$

In other words, there is a set of more than e rows of M' at which $M'|_I$ is all ones, $M'|_S$ has weight exactly u , and $M'|_Z$ is all zeros, as required by Definition 13. $\square$

The special case $u = g + 1$ in the above proposition is particularly interesting. A chain of reductions between strongly disjunct and threshold disjunct matrices in this case implied by the above result is schematically shown below.

$$\begin{array}{c} (2d, e; g + 1, g)\text{-disjunct} \\ \downarrow \\ \text{strongly } (2d, e; g + 1)\text{-disjunct} \\ \downarrow \\ (d, e; g + 1, g)\text{-disjunct} \\ \downarrow \\ \text{strongly } (d, e; g + 1)\text{-disjunct} \end{array}$$

Therefore, when the upper threshold u is more than the gap parameter g by one (equivalently, when the lower threshold ℓ is one), the two notions of threshold disjunct matrices and strongly disjunct matrices become equivalent up to a multiplicative factor in the sparsity parameter d . As discussed in Section 2, almost matching lower bounds and upper bounds on the number of rows m achievable by a strongly $(d, e; g + 1)$ -disjunct matrix are known. Asymptotically, the number of rows must always satisfy $m = \Omega(d^{g+2} \log_d n + ed^{g+1})$ and moreover, a probabilistic construction achieves $m = O(d^{g+2} \log(n/d))$ and $e = \Omega(d \log(n/d))$ with probability $1 - o(1)$ (see Table 1). As

a result, the upper and lower bounds on the number of rows of strongly disjunct and threshold disjunct matrices become equivalent up to multiplicative constants when the lower threshold is one.

Proposition 15 asserts that the notion of strongly disjunct matrices is in general stronger than threshold disjunct matrices. As we will see below, the former becomes strictly stronger when $\ell > 1$. As the lower threshold ℓ becomes larger, the discrepancy between the number of rows achievable by threshold disjunct matrices and strongly disjunct matrices becomes more significant (see Table 1).

Observation 3

As pointed out after Definition 13, the generalized definition of regular matrices may affect the bounds obtained by our probabilistic and explicit constructions (Constructions 2 and 4) only by hidden factors depending on u (essentially without any change in the proofs). For the case of generalized disjunct matrices, however, the bounds may substantially change depending on the gap parameter g .

Below we generalize Lemma 8 for the case of threshold-disjunct matrices and show that Construction 2 results in a $(d, \Omega_u(pd \log(n/d)/(1-p)^2); u, g)$ -disjunct matrix (with probability $1 - o(1)$) if the number of measurements is increased by a factor $O(d^g)$. More precisely, we can now show the following lemma.

Lemma 16 (Lemma 8, generalized). *For every $p \in [0, 1)$ and integer parameters $u > g \geq 0$, Construction 2 with $m' = O_u(d^{g+2} \log(n/d)/(1-p)^2)$ outputs a $(d, \Omega_u(pm'/d^{g+1}); u, g)$ -disjunct matrix with probability $1 - o(1)$.*

Proof. The proof essentially follows along the same lines as the proof of Lemma 8. The difference, compared to the case $g = 0$ covered by Lemma 8, is that we have a set I of distinguished columns $I \subseteq [n]$ in Definition 13 where $|I| = g + 1$ and the random vector $\mathbf{w}$ in the proof of Lemma 8 must have ones at all positions picked by I . With this requirement, the lower bound on the success probability q in (3) becomes $c = \Omega_u(1/d^{g+1})$. The rest of the proof remains unchanged except for the new lower bound on c , which makes the error tolerance parameter e in the proof lower bounded by $\Omega(pm'/d^{g+1})$, while increasing the parameter m' to a quantity upper bounded by $O_u(d^{g+2} \log(n/d)/(1-p)^2)$. $\square$

Observation 4

Lemma 7 can be extended to positive gaps as follows.

Lemma 17 (Lemma 7, generalized). *Suppose that M_1 is a $(d, e_1; u, g + 1)$ -regular and M_2 is a strongly $(2d, e_2; g + 1)$ -disjunct matrix. Then $M_1 \odot M_2$ is a $(d, (e_1 + 1)(e_2 + 1) - 1; u, g)$ -disjunct matrix where $u := \ell + g$.*

Proof. Let $M := M_1 \odot M_2$. Towards verifying that M satisfies the requirements of Definition 13, consider a set $I \subseteq [n]$ of distinguished columns of M , where n is the number of columns of the matrices and $|I| = g + 1$, in addition to critical and zero sets $S, Z \subseteq [n]$ as in Definition 13 satisfying $|Z| \leq |S|$. Index the rows of M naturally by the elements of $[m_1] \times [m_2]$, where m_1 and m_2 are the number of rows of M_1 and M_2 , respectively, and the (i, j) th row of M is the bitwise disjunction of the i th row of M_1 and the j th row of M_2 .

Let $S' := S \setminus I$ and $Z' := Z \cup (S \setminus I)$. Observe that $|Z| \leq |S| \leq |S'| + g + 1 = |S| + |I|$ and $|Z'| \leq 2d$. From Definition 13, there is a set $E_1 \subseteq [m_1]$ of size more than e_1 such that $M_1|_{S'}$ has weight exactly $\ell - 1 = u - |I|$, and $M_2|_Z$ is all zeros. Moreover, there is a set $E_2 \subseteq [m_2]$ of size more than e_2 at which $M_2|_I$ has all ones and $M_2|_{Z'}$ has all zeros. This means that, at all rows

corresponding to $E_1 \times E_2$, the product matrix M has weight exactly $\ell - 1 + |I| = u$ at positions corresponding to S and all zeros at positions corresponding to Z . Therefore, M indeed u -satisfies any choice of the sets I, S, Z at more than $(e_1 + 1)(e_2 + 1) - 1$ rows. $\square$

Consequently, using the coding-theoretic construction of strongly disjunct matrices described in Appendix A, our explicit constructions of $(d, e; u)$ -disjunct matrices obtained in Section 3 can be extended to the positive gap model at the cost of a factor $O(d^g)$ increase in the number of measurements. The results from combining the above lemma with various constructions of regular and strongly disjunct matrices are summarized in Table 1.

5 Concluding remarks

In this work we have introduced the combinatorial notion of regular binary matrices, that is used as an intermediate tool towards obtaining threshold testing designs.

Even though our construction, assuming an optimal lossless condenser, matches the probabilistic upper bound for regular matrices, the number of measurements in the resulting threshold testing scheme (obtained from the simple direct product in Construction 1) becomes larger than the probabilistic upper bound by a factor of $\Omega(d \log n)$. Thus, an outstanding question is coming up with a direct construction of disjunct matrices that matches the probabilistic upper bound. Despite this, the notion of regular matrices may be of independent interest, and an interesting question is to obtain (nontrivial) concrete lower bounds on the number of rows of such matrices in terms of the parameters n, d, e, u (and the gap parameter g in the generalized definition of Section 4).

Moreover, in this work we have assumed the upper threshold u to be a fixed constant, allowing the constants hidden in asymptotic notions to have a poor dependence on u . An outstanding question is whether the number of measurements can be reasonably controlled when the upper threshold u and possibly the gap parameter g become large; e.g., $g, u = \Omega(d)$.

Another interesting problem is decoding. While our constructions can combinatorially guarantee identification of sparse vectors, for applications it is important to have an efficient reconstruction algorithm as well. Contrary to the case of strongly disjunct matrices that allow a straightforward decoding procedure (cf. [6]), it is not clear whether in general our notion of disjunct matrices allow efficient decoding, and thus it becomes important to look for constructions that are equipped with efficient reconstruction algorithms.

Finally, for clarity of the exposition, in this work we have only focused on asymptotic trade-offs, and it would be nice to obtain good, finite length, estimates on the obtained bounds that are useful for applications.

References

- [1] A. Ben-Aroya and A. Ta-Shma, *Constructing small-bias sets from algebraic-geometric codes*, Proceedings of the 50th annual IEEE symposium on foundations of computer science (FOCS), 2009.
- [2] A. Blass and Y. Gurevich, *Pairwise testing*, Bulletin of the EATCS **78** (2002), 100–132.
- [3] W.J. Bruno, E. Knill, D.J. Balding, D.C. Bruce, N.A. Doggett, W.W. Sawhill, R.L. Stallings, C.C. Whittaker, and D.C. Torney, *Efficient pooling designs for library screening*, Genomics **26** (1995), no. 1, 21–30.
- [4] M. Capalbo, O. Reingold, S. Vadhan, and A. Wigderson, *Randomness conductors and constant-degree expansion beyond the degree/2 barrier*, Proceedings of the 34th annual ACM symposium on theory of computing (STOC), 2002, pp. 659–668.
- [5] H-B. Chen, D-Z. Du, and F-K. Hwang, *An unexpected meeting of four seemingly unrelated problems: graph testing, DNA complex screening, superimposed codes and secure key distribution*, Journal of Combinatorial Optimization **14** (2007), no. 2-3, 121–129.

- [6] H-B. Chen and H-L. Fu, *Nonadaptive algorithms for threshold group testing*, Discrete Applied Mathematics **157** (2009), 1581–1585.
- [7] H-B. Chen, H-L. Fu, and F-K. Hwang, *An upper bound of the number of tests in pooling designs for the error-tolerant complex model*, Optimization Letters **2** (2008), no. 3, 425–431.
- [8] Y. Cheng and D-Z. Du, *New constructions of one- and two-stage pooling designs*, Journal of Computational Biology **15** (2008), no. 2, 195–205.
- [9] M. Cheraghchi, *Noise-resilient group testing: Limitations and constructions*, Proceedings of the 17th international symposium on fundamentals of computation theory (FCT), 2009, pp. 62–73.
- [10] ———, *Improved constructions for non-adaptive threshold group testing*, Proceedings of the 37th international colloquium on automata, languages and programming (ICALP), 2010. arXiv: cs.DM/1002.2244.
- [11] R. Clifford, K. Efremenko, E. Porat, and A. Rothschild, *k-mismatch with don't cares*, Proceedings of the 15th european symposium on algorithm (ESA), 2007, pp. 151–162.
- [12] G. Cormode and S. Muthukrishnan, *What's hot and what's not: tracking most frequent items dynamically*, ACM Transactions on Database Systems **30** (2005), no. 1, 249–278.
- [13] ———, *Combinatorial algorithms for compressed sensing*, Proceedings of information sciences and systems, 2006, pp. 198–201.
- [14] P. Damaschke, *Threshold group testing*, General theory of information transfer and combinatorics, 2006, pp. 707–718.
- [15] R. Dorfman, *The detection of defective members of large populations*, Annals of Mathematical Statistics **14** (1943), 436–440.
- [16] D-Z. Du and F. Hwang, *Combinatorial group testing and its applications*, Second, World Scientific, 2000.
- [17] D-Z. Du and F-K. Hwang, *Pooling designs and nonadaptive group testing*, World Scientific, 2006.
- [18] A. D'yachkov, P. Vilenkin, A. Macula, and D. Torney, *Families of finite sets in which no intersection of ℓ sets is covered by the union of s others*, Journal of Combinatorial Theory, Series A **99** (2002), 195–218.
- [19] M. Farach, S. Kannan, E. Knill, and S. Muthukrishnan, *Group testing problems with sequences in experimental molecular biology*, Proceedings of compression and complexity of sequences, 1997, pp. 357–367.
- [20] H. Gao, F.K. Hwang, M.T. Thai, W. Wu, and T. Znati, *Construction of $d(H)$ -disjunct matrix for group testing in hypergraphs*, Journal of Combinatorial Optimization **12** (2006), 297–301.
- [21] A. Garcia and H. Stichtenoth, *A tower of Artin-Schreier extensions of function fields attaining the Drinfeld-Vlăduț bound*, Invent. Math. **121** (1995), 211–222.
- [22] V. Guruswami, C. Umans, and S. Vadhan, *Unbalanced expanders and randomness extractors from Parvaresh-Vardy codes*, Proceedings of the 22nd IEEE conference on computational complexity (CCC), 2007.
- [23] E-S. Hong and R.E. Ladner, *Group testing for image compression*, Data compression conference, 2000, pp. 3–12.
- [24] W.H. Kautz and R.C. Singleton, *Nonrandom binary superimposed codes*, IEEE Transactions on Information Theory **10** (1964), 363–377.
- [25] H.K. Kim and V. Lebedev, *On optimal superimposed codes*, Journal of Combinatorial Designs **12** (2004), 79–91.
- [26] J. H. van Lint, *Introduction to coding theory*, Third, Graduate Texts in Mathematics, vol. 86, Springer Verlag, 1998.
- [27] A.J. Macula, *Probabilistic nonadaptive group testing in the presence of errors and DNA library screening*, Annals of Combinatorics **3** (1999), no. 1, 61–69.
- [28] F.J. MacWilliams and N.J. Sloane, *The theory of error-correcting codes*, North Holland, 1977.
- [29] H-Q. Ngo and D-Z. Du, *A survey on combinatorial group testing algorithms with applications to DNA library screening*, DIMACS Series on Discrete Math. and Theoretical Computer Science **55** (2000), 171–182.
- [30] E. Porat and A. Rothschild, *Explicit non-adaptive combinatorial group testing schemes*, Proceedings of the 35th international colloquium on automata, languages and programming (ICALP), 2008, pp. 748–759.
- [31] J. Radhakrishnan and A. Ta-Shma, *Tight bounds for depth-two superconcentrators*, Proceedings of the 38th annual IEEE symposium on foundations of computer science (FOCS), 1997, pp. 585–594.
- [32] R.M. Roth, *Introduction to coding theory*, Cambridge University Press, 2006.
- [33] A. Schliep, D. Torney, and S. Rahmann, *Group testing with DNA chips: Generating designs and decoding experiments*, Proceedings of computational systems bioinformatics, 2003.

- [34] D.R. Stinson and R. Wei, *Generalized cover-free families*, Discrete Mathematics **279** (2004), 463–477.
- [35] D.R. Stinson, R. Wei, and L. Zhu, *Some new bounds for cover-free families*, Journal of Combinatorial Theory, Series A **90** (2000), 224–234.
- [36] M.A. Tsfasman, S.G. Vlăduț, and Th. Zink, *Modular curves, Shimura curves, and Goppa codes better than the Varshamov-Gilbert bound*, Mathematische Nachrichten **109** (1982), 21–28.
- [37] J. Wolf, *Born-again group testing: multiaccess communications*, IEEE Transactions on Information Theory **31** (1985), 185–191.
- [38] W. Wu, Y. Huang, X. Huang, and Y. Li, *On error-tolerant DNA screening*, Discrete Applied Mathematics **154** (2006), no. 12, 1753–1758.
- [39] W. Wu, Y. Li, C.H. Huang, and D.Z. Du, *Molecular biology and pooling design*, Data Mining in Biomedicine **7** (2008), 133–139.

A Strongly disjoint matrices from codes

In this appendix we describe a construction of strongly disjoint matrices (as in Definition 3) which is a straightforward extension of the classical result of Kautz and Singleton [24] for construction of combinatorial designs. Construction 5 explains the idea, which is analyzed in Lemma 18 below. In this section we use standard tools from the theory of error-correcting codes. The interested reader is referred the standard texts in coding theory (e.g., the books by MacWilliams and Sloane [28], van Lint [26], and Roth [32]) for background.

- *Given:* An $(\tilde{n}, k, \tilde{d})_q$ error-correcting code¹⁰ $\mathcal{C} \subseteq [q]^{\tilde{n}}$, and integer parameter $u > 0$.
- *Output:* An $m \times n$ Boolean matrix M , where $n = q^k$, and $m = \tilde{n}q^u$.
- *Construction:* First, consider the mapping $\varphi: [q] \rightarrow \{0, 1\}^{q^u}$ from q -ary symbols to column vectors of length q^u defined as follows. Index the coordinates of the output vector by the u -tuples from the set $[q]^u$. Then $\varphi(x)$ has a 1 at position $(a_1, \dots, a_u)$ if and only if there is an $i \in [u]$ such that $a_i = x$. Arrange all codewords of $\mathcal{C}$ as columns of an $\tilde{n} \times q^k$ matrix M' with entries from $[q]$. Then replace each entry x of M' with $\varphi(x)$ to obtain the output $m \times n$ matrix M .

Construction 5: Extension of Kautz-Singleton’s method [24].

Lemma 18. *Construction 5 outputs a strongly $(d, e; u)$ -disjoint matrix for every $d < (\tilde{n} - e)/((\tilde{n} - \tilde{d})u)$.*

Proof. Let $C := \{c_1, \dots, c_u\} \subseteq [n]$ and $C' := \{c'_1, \dots, c'_d\} \subseteq [n]$ be disjoint subsets of column indices. We wish to show that, for more than e rows of M , the entries at positions picked by C are all-ones while those picked by C' are all-zeros. For each $j \in [n]$, denote the j th column of M' by $M'(j)$, and let $M'(C) := \{M'(c_j) : j \in [u]\}$, and $M'(C') := \{M'(c'_j) : j \in [d]\}$.

From the minimum distance of $\mathcal{C}$, we know that every two distinct columns of M' agree in at most $\tilde{n} - \tilde{d}$ positions. By a union bound, for each $i \in [d]$, the number of positions where $M'(c'_i)$ agrees with one or more of the codewords in $M'(C)$ is at most $u(\tilde{n} - \tilde{d})$, and the number of positions where some vector in $M'(C')$ agrees with one or more of those in $M'(C)$ is at most $du(\tilde{n} - \tilde{d})$.

By assumption, we have $\tilde{n} - du(\tilde{n} - \tilde{d}) > e$, and thus, for a set $E \subseteq [\tilde{n}]$ of size greater than e , at positions picked by E none of the codewords in $M'(C')$ agree with any of the codewords in $M'(C)$.

¹⁰We use the notation $(\tilde{n}, k, \tilde{d})_q$ code for a q -ary code of length $\tilde{n}$, size q^k , and minimum distance at least $\tilde{d}$.

Now let $w \in [q]^n$ be any of the rows of M' picked by E , and consider the $q^u \times n$ Boolean matrix W formed by applying the mapping $\varphi(\cdot)$ on each entry of w . We know that $\{w(c_j) : j \in [u]\} \cap \{w(c'_j) : j \in [d]\} = \emptyset$. Thus we observe that the particular row of W indexed by $(w(c_1), \dots, w(c_u))$ (and in fact, any of its permutations) must have all-ones at positions picked by C and all-zeros at those picked by C' . As any such row is a distinct row of M , it follows that M is strongly $(d, e; u)$ -disjunct. $\square$

Here we mention a few specific instantiations of the above construction. Namely, we will first consider the family of Reed-Solomon codes, that are also used in the original work of Kautz and Singleton [24], and then move on to the family of algebraic geometric (AG) codes on the the Tsfasman-Vlăduț-Zink (TVZ) bound, Hermitian codes, and finally, codes on the Gilbert-Varshamov (GV) bound.

Reed-Solomon codes: Let $p \in [0, 1)$ be an arbitrary “noise” parameter. If we take $\mathcal{C}$ to be an $[\tilde{n}, k, \tilde{d}]_{\tilde{n}}$ Reed-Solomon code over an alphabet of size $\tilde{n}$ (which we assume to be a prime power), where $\tilde{d} = \tilde{n} - k + 1$, we get a strongly disjunct $(d, e; u)$ -matrix with $m = O(du \log n / (1 - p))^{u+1}$ rows and $e = p\tilde{n} = \Omega(pdu(\log n)/(1 - p))$.

AG codes on the TVZ bound: Another interesting family for the code $\mathcal{C}$ is the family of algebraic geometric codes that attain the Tsfasman-Vlăduț-Zink bound (cf. [21, 36]). This family is defined over any alphabet size $q \geq 49$ that is a square prime power, and achieves a minimum distance $\tilde{d} \geq \tilde{n} - k - \tilde{n}/(\sqrt{q} - 1)$. Let $e := pn$, for a noise parameter $p \in [0, 1)$. By Lemma 18, the underlying code $\mathcal{C}$ needs to have minimum distance at least $\tilde{n}(1 - (1 - p)/(du))$. Thus in order to be able to use the above-mentioned family of AG codes, we need to have $q \gg (du/(1 - p))^2 =: q_0$. Let us take an appropriate $q \in [2q_0, 8q_0]$, and following Lemma 18, $\tilde{n} - \tilde{d} = \lceil \tilde{n}(1 - p)/(du) \rceil$. Thus, the dimension of $\mathcal{C}$ becomes at least

$$k \geq \tilde{n} - \tilde{d} - \frac{\tilde{n}}{\sqrt{q} - 1} = \Omega\left(\frac{\tilde{n}(1 - p)}{du}\right) = \Omega(\tilde{n}/\sqrt{q_0}),$$

and subsequently¹¹ we get that $\log n = k \log q \geq k = \Omega(\tilde{n}/\sqrt{q_0})$. Now, noting that $m = q^u \tilde{n}$, we conclude that

$$m = q^u \tilde{n} = O(q_0^{u+1/2} \log n) = O\left(\frac{du}{1 - p}\right)^{2u+1} \log n,$$

and $e = \Omega(pdu(\log n)/(1 - p))$.

We see that the dependence of the number of measurements on the sparsity parameter d is worse for AG codes than Reed-Solomon codes by a factor d^u , but the construction from AG codes benefits from a linear dependence on $\log n$, compared to $\log^{u+1} n$ for Reed-Solomon codes. Thus, AG codes become more favorable only when the sparsity is substantially low; namely, when $d \ll \log n$.

Hermitian codes. A particularly nice family of AG codes arises from the Hermitian function field. Let q' be a prime power and $q := q'^2$. Then the Hermitian function field over $\mathbb{F}_q$ is a finite extension of the rational function field $\mathbb{F}_q(x)$, denoted by $\mathbb{F}_q(x, y)$, where we have $y^{q'} + y = x^{q'+1}$. The structure of this function field is relatively well understood and the family of Goppa codes defined over the rational points of the Hermitian function field is known as Hermitian codes. This family is recently used by Ben-Aroya and Ta-Shma [1] for construction of small-bias sets. Below we quote some parameters of Hermitian codes from their work.

The number of rational points of the Hermitian function field is equal to $q'^3 + 1$, which includes a common pole Q_∞ of x and y . The genus of the function field is $\tilde{g} = q'(q' - 1)/2$. For some integer

¹¹Note that, given the parameters p, d, n , the choice of q depends on p, d , as explained above, and then one can choose the code length $\tilde{n}$ to be the smallest integer for which we have $q^k \geq n$. But for the sake of clarity we have assumed that $q^k = n$, which does not affect the asymptotic bounds.

parameter r , we take $G := rQ_\infty$ as the divisor defining the Riemann-Roch space $\mathcal{L}(G)$ of the code $\mathcal{C}$, and the set of rational points except Q_∞ as the evaluation points of the code. Thus the length of $\mathcal{C}$ becomes $\tilde{n} = q'^3$. Moreover, the minimum distance of the code is $\tilde{d} = n - \deg(G) = n - r$. When $r \geq 2\tilde{g} - 1$, the dimension of the code is given by the Riemann-Roch theorem, which is equal to $r - \tilde{g} + 1$. For the low-degree regime where $r < 2\tilde{g} - 1$, the dimension k of the code is the size of the Wirestraussem semigroup of G , which turns out to be the set $W = \{(i, j) \in \mathbb{N}^2 : j \leq q' - 1 \wedge iq' + j(q' + 1) \leq r\}$.

Now, given parameters d, p of the disjunct matrix, define $\rho := (1 - p)/((d + 1)u)$, take the alphabet size q as a square prime power, and set $r := \rho q^{3/2}$. First we consider the case where $r < 2\tilde{g} - 1 = 2q - 2\sqrt{q} - 1$. In this case, the dimension of the Hermitian code becomes $k = |W| = \Omega(r^2/q) = \Omega(\rho^2 q^2)$. The distance $\tilde{d}$ of the code satisfies $\tilde{d} = \tilde{n} - r \geq \tilde{n}(1 - \rho)$ and thus, for $e := p\tilde{n}$, conditions of Lemma 18 are satisfied. The number of the rows of the resulting measurement matrix becomes $m = q^{u+3/2}$, and we have $n = q^k$. Therefore,

$$\log n = k \log q \geq k = \Omega(\rho^2 q^2) \Rightarrow q = O(\sqrt{\log n}/\rho) \Rightarrow m = O\left(\left(\frac{d\sqrt{\log n}}{1-p}\right)^{u+3/2}\right),$$

and in order to ensure that $r < 2\tilde{g} - 1$, we need to have $du/(1 - p) \gg \sqrt{\log n}$. On the other hand, when $du/(1 - p) \ll \sqrt{\log n}$, we are in the high-degree regime, in which case the dimension of the code becomes $k = r - \tilde{g} + 1 = \Omega(r) = \Omega(\rho q^{3/2})$, and we will thus have

$$q = O((\log n/\rho)^{2/3}) \Rightarrow m = O\left(\left(\frac{d \log n}{1-p}\right)^{1+2u/3}\right)$$

Altogether, we conclude that Construction 5 with Hermitian codes results in a strongly $(d, e; u)$ -disjunct matrix with

$$m = O\left(\left(\frac{d\sqrt{\log n}}{1-p} + \left(\frac{d \log n}{1-p}\right)^{2/3}\right)^{u+3/2}\right)$$

rows, where $e = p \cdot \Omega(d \log n/(1 - p) + (d\sqrt{\log n}/(1 - p))^{3/2})$. Compared to the Reed-Solomon codes, the number of measurements has a slightly worse dependence on d , but a much better dependence on n . Compared to AG codes on the TVZ bound, the dependence on d is better while the dependence on n is inferior.

Codes on the GV bound. A q -ary $(\tilde{n}, k, \tilde{d})$ -code (of sufficiently large length) is said to be on the Gilbert-Varshamov bound if it satisfies $k \geq \tilde{n}(1 - h_q(\tilde{d}/\tilde{n}))$, where $h_q(\cdot)$ is the q -ary entropy function defined as

$$h_q(x) := x \log_q(q - 1) - x \log_q(x) - (1 - x) \log_q(1 - x).$$

It is well known that a random linear code achieves the bound with overwhelming probability (cf. [28]). Now we apply Lemma 18 on a code on the GV bound, and calculate the resulting parameters. Let $\rho := (1 - p)/(4du)$, choose any alphabet size $q \in [1/\rho, 2/\rho]$, and let $\mathcal{C}$ be any q -ary code of length $\tilde{n}$ on the GV bound, with minimum distance $\tilde{d} \geq \tilde{n}(1 - 2/q)$. By the Taylor expansion of the function $h_q(x)$ around $x = 1 - 1/q$, we see that the dimension of $\mathcal{C}$ asymptotically behaves as $k = \Theta(\tilde{n}/(q \log q))$. Thus, the number of columns of the resulting measurement matrix becomes $n = q^k = 2^{\Omega(\tilde{n}/q)}$. Moreover, the number m of its rows becomes

$$m = q^u \tilde{n} = O(q^{u+1} \log n) = O((d/(1 - p))^{u+1} \log n),$$

and the matrix becomes strongly $(d, e; u)$ -disjunct for $e = p\tilde{n} = \Omega(pd \log n)/(1 - p)$.

We remark that for the range of parameters that we are interested in, Porat and Rothschild [30] have come up with a deterministic construction of linear codes on the GV bound that runs in time $\text{poly}(q^k)$ (and thus, polynomial in the size of the resulting measurement matrix).

Their construction is based on a derandomization of the probabilistic argument for random linear codes using the method of conditional expectations, and as such, can be considered *weakly explicit* (in the sense that, the entire measurement matrix can be computed in polynomial time in its length; whereas for a fully explicit construction one must ideally be able to deterministically compute any single entry of the measurement matrix in time $\text{poly}(d, \log n)$, which is not the case for this construction). Altogether, we obtain the following result.

Theorem 19. *There is an algorithm that, given integer parameters $d \leq n$ and $u > 0$ and real parameter $p \in [0, 1)$ outputs an $m \times n$ binary matrix which is strongly $(d, e; u)$ -disjunct. The parameters m and e satisfy the bounds $m = O((d/(1-p))^{u+1} \log n)$ and $e = \Omega(pd(\log n)/(1-p))$. Moreover, the running time of the algorithm is polynomial in mn . $\square$*

Using a standard probabilistic argument it is easy to see that a random $m \times n$ matrix, where each entry is an independent Bernoulli random variable with probability $1/d$ of being 1, is with overwhelming probability strongly $(d, e; u)$ -disjunct for $e = \Omega(pd \log(n/d)/(1-p)^2)$ and $m = O(d^{u+1}(\log(n/d))/(1-p)^2)$ (the proof is very similar to the proof of Lemma 8). Thus we see that, for a fixed p , Construction 5 when using codes on the GV bound almost matches these parameters. Moreover, the explicit construction based on Reed-Solomon codes possesses the “right” dependence on the sparsity d , while AG codes on the TVZ bound have a matching dependence on the vector length n with random measurement matrices, and finally, the trade-off offered by the construction based on Hermitian codes lies in between the one for Reed-Solomon codes and AG codes.